

Fermion-Boson-Type Subquark Model and $\Delta F = 2$ Phenomena

TAKEO MATSUSHIMA *

*17-1 1-Chome Azura Ichinomiya,
Aichi-Prefecture 491, Japan*

Abstract

$\Delta F = 2$ phenomena in all neutral pseudo scalar meson systems are investigated in the context of the fermion-boson-type subquark model. In this model the mass difference between the heavier neutral pseudo scalar meson and the lighter one and indirect CP violation associated with the mixing are unifyingly explained by the neutral scalar subquark exchanges between two quarks inside the present meson. We obtain the mass differences : $\Delta M_D \approx 10^{-14}$ GeV, $\Delta M_{B_s} \approx 10^{-11}$ GeV, $\Delta M_{T_u} \approx 10^{(-10 \sim -9)}$ GeV and $\Delta M_{T_c} \approx 10^{(-8 \sim -7)}$ GeV. It is also predicted that indirect CP violations in D- and B-meson systems are same as in K-meson system.

*† Available mail address is as follows :

Nagoya University Department of Physics Chikusa-ku Nagoya 464-01 Japan
e-mail : mtakeo@eken.phys.nagoya-u.ac.jp

1 Introduction

Since the first observation of CP violation in the K^0 system in 1964[1], this evidence has not been explained in a successful scenario. It was soon after this experiment in the same year that Wolfenstein proposed the idea of super-weak theory that CP violation occurs only in the effective Hamiltonian which describes the time evolution of the K^0 system and its strength is 10^{-9} times smaller than the standard weak interactions. In 1973 Kobayashi and Maskawa introduced the 3×3 unitary matrix which mixes three generations in the context of the electroweak unified gauge theory called the standard model (SM) usually. This model has the ability to explain CP violations not only in the effective Hamiltonian of the second order weak interaction but also in the decay amplitudes if two different decay processes contribute to one final state. In the latter case the parameter $Re(\epsilon'/\epsilon)_K \neq 0$ is expected although a large top quark mass effect gives the prediction of very small value[3]. On the other hand the super-weak theory predict the exact zero value. The experimental efforts about $Re(\epsilon'/\epsilon)_K$ so far have not given a conclusive answer. Recently CP violations in heavier neutral mesons (D^0 , B^0) have been discussed as the concerning experiments were carried out (though not yet been observed) and in the near future the real features will become clear. So the totally understandable scenario of CP violation phenomena is required. The mass difference (ΔM_P) between heavier neutral pseudo scalar (P -) meson and lighter one is considered to be essentially connected with CP violations because the origin of both phenomena comes from the off diagonal matrix elements of the mass matrix (M_{ij} , $i, j = 1, 2$) and the decay matrix (Γ_{ij} , $i, j = 1, 2$). Comparing with CP violations, the experiments of ΔM_P are a little abundant, e.g., there are ΔM_K , ΔM_{B_d} , the upper bound of ΔM_D , and the lower bound of ΔM_{B_s} [4]. Theoretical analyses about them are roughly in two ways, e.g., the estimation of M_{12} by the superweak theory or the box diagram calculation in the SM with (or without) long distance contributions. Therefore the aim of the present stage is to clarify what kind of dynamics controls M_{ij} and Γ_{ij} . In this paper we investigate this issue in the context of the Fermion-Boson-type subquark model (FB-model) inspired by some type of gauge theory which proposed by the author[5]. In this model the assumption that the “neutral scalar subquarks ($\mathbf{y}$) exchange” between two quarks inside the present P -meson plays the essential role. So this scenario may be said a realization of Wolfenstein’s super-weak idea[2]. In Sect.2 we mention the gauge theory which inspires quark-lepton composite scenario. In Sect.3 we review the FB-model by which we study the present issue. In Sect.4 we investigate the mass

differences and CP violations in neutral pseudo scalar meson systems by using this FB-model.

2 Gauge theory inspiring quark-lepton composite scenario

In our model the existence of fundamental matter fields (preon) are inspired by the gauge theory with Cartan connections[5]. Let us briefly summarize the basic features of that. Generally gauge fields, including gravity, are considered as geometrical objects, that is, connection coefficients of principal fiber bundles. It is said that there exist some different points between Yang-Mills gauge theories and gravity, though both theories commonly possess fiber bundle structures. The latter has the fiber bundle related essentially to 4-dimensional space-time freedoms but the former is given, in an ad hoc way, the one with the internal space which has nothing to do with the space-time coordinates. In case of gravity it is usually considered that there exist ten gauge fields, that is, six spin connection fields in $SO(1, 3)$ gauge group and four vierbein fields in $GL(4, R)$ gauge group from which the metric tensor $g^{\mu\nu}$ is constructed in a bilinear function of them. Both altogether belong to Poincaré group $ISO(1, 3) = SO(1, 3) \otimes R^4$ which is semi-direct product. In this scheme spin connection fields and vierbein fields are independent but only if there is no torsion, both come to have some relationship. Seeing this, $ISO(1, 3)$ gauge group theory has the logical weak point not to answer how two kinds of gravity fields are related to each other intrinsically.

In the theory of Differential Geometry, S.Kobayashi has investigated the theory of “Cartan connection”[6]. This theory, in fact, has ability to reinforce the above weak point. The brief recapitulation is as follows. Let $E(B_n, F, G, P)$ be a fiber bundle (which we call Cartan-type bundle) associated with a principal fiber bundle $P(B_n, G)$ where B_n is a base manifold with dimension “ n ”, G is a structure group, F is a fiber space which is homogeneous and diffeomorphic with G/G' where G' is a subgroup of G . Let $P' = P'(B_n, G')$ be a principal fiber bundle, then P' is a subbundle of P . Here let it be possible to decompose the Lie algebra $\mathfrak{g}$ of G into the subalgebra $\mathfrak{g}'$ of G' and a vector space $\mathfrak{f}$ such as :

$$\mathfrak{g} = \mathfrak{g}' + \mathfrak{f}, \quad \mathfrak{g}' \cap \mathfrak{f} = 0, \quad (1)$$

$$[\mathfrak{g}', \mathfrak{g}'] \subset \mathfrak{g}', \quad (2)$$

$$[\mathbf{g}', \mathbf{f}] \subset \mathbf{f}, \quad (3)$$

$$[\mathbf{f}, \mathbf{f}] \subset \mathbf{g}', \quad (4)$$

where $\dim \mathbf{f} = \dim F = \dim G - \dim G' = \dim B_n = n$. The homogeneous space $F = G/G'$ is said to be “weakly reductive” if there exists a vector space $\mathbf{f}$ satisfying Eq.(1) and (3). Further F satisfying Eq(4) is called “symmetric space”. Let ω denote the connection form of P and $\bar{\omega}$ be the restriction of ω to P' . Then $\bar{\omega}$ is a $\mathbf{g}$ -valued linear differential 1-form and we have :

$$\omega = g^{-1}\bar{\omega}g + g^{-1}dg, \quad (5)$$

where $g \in G$, $dg \in T_g(G)$. ω is called the form of “Cartan connection” in P .

Let the homogeneous space $F = G/G'$ be weakly reductive. The tangent space $T_O(F)$ at $o \in F$ is isomorphic with $\mathbf{f}$ and then $T_O(F)$ can be identified with $\mathbf{f}$ and also there exists a linear $\mathbf{f}$ -valued differential 1-form(denoted by θ) which we call the “form of soldering”. Let ω' denote a $\mathbf{g}'$ -valued 1-form in P' , we have :

$$\bar{\omega} = \omega' + \theta. \quad (6)$$

The dimension of vector space $\mathbf{f}$ and the dimension of base manifold B_n is the same “ n ”, and then $\mathbf{f}$ can be identified with the tangent space of B_n at the same point in B_n and θ s work as n -bein fields. In this case ω' and θ unifyingly belong to group G . Here let us call such a mechanism “Soldering Mechanism”.

Drechsler has found out the useful aspects of this theory and investigated a gravitational gauge theory based on the concept of the Cartan-type bundle equipped with the Soldering Mechanism[7]. He considered $F = SO(1, 4)/SO(1, 3)$ model. Homogeneous space F with $\dim = 4$ solders 4-dimensional real space-time. The Lie algebra of $SO(1, 4)$ corresponds to $\mathbf{g}$ in Eq.(1), that of $SO(1, 3)$ corresponds to $\mathbf{g}'$ and $\mathbf{f}$ is 4-dimensional vector space. The 6-dimensional spin connection fields are $\mathbf{g}'$ -valued objects and vierbein fields are $\mathbf{f}$ -valued, both of which are unified into the members of $SO(1, 4)$ gauge group. We can make the metric tensor $\mathbf{g}^{\mu\nu}$ as a bilinear function of $\mathbf{f}$ -valued vierbein fields. Inheriting Drechsler’s study the author has investigated the quantum theory of gravity[5]. The key point for this purpose is that F is a symmetric space because $\mathbf{f}$ s are satisfied with Eq.(4). Using this symmetric nature we can inquire into making a quantum gauge theory, that is, constructing $\mathbf{g}'$ -valued Faddeev-Popov ghost, anti-ghost, gauge fixing, gaugeon and its pair field as composite fusion fields of $\mathbf{f}$ -valued gauge fields by use of Eq.(4) and also naturally inducing BRS-invariance.

Comparing such a scheme of gravity, let us consider Yang-Mills gauge theories. Usually when we make the Lagrangian density $\mathcal{L} = \text{tr}(\mathbf{F} \wedge \mathbf{F}^*)$ ($\mathbf{F}$ is a field strength), we must borrow a metric tensor $\mathbf{g}^{\mu\nu}$ from gravity to get $\mathbf{F}^*$ and also for Yang-Mills gauge fields to propagate in the 4-dimensional real space-time. This seems to mean that “there is a hierarchy between gravity and other three gauge fields (electromagnetic, strong, and weak)”. But is it really the case? As an alternative thought we can think that all kinds of gauge fields are “equal”. Then it would be natural for the question “What kind of equality is that?” to arise. In other words, it is the question that “What is the minimum structure of the gauge mechanism which four kinds of forces are commonly equipped with?”. For answering this question, let us make an assumption: “*Gauge fields are Cartan connections equipped with Soldering Mechanism.*” In this meaning all gauge fields are equal. If it is the case three gauge fields except for gravity are also able to have their own metric tensors and to propagate in the real space-time without the help of gravity. Such a model has already been investigated in Ref.[5].

Let us discuss them briefly. It is found that there are four types of sets of classical groups with smaller dimensions which admit Eq.(1,2,3,4), that is, $F = SO(1, 4)/SO(1, 3)$, $SU(3)/U(2)$, $SL(2, C)/GL(1, C)$ and $SO(5)/SO(4)$ with $\dim F = 4$ [8]. Note that the quality of “ $\dim 4$ ” is very important because it guarantees F to solder to 4-dimensional real space-time and all gauge fields to work in it. The model of $F = SO(1, 4)/SO(1, 3)$ for gravity is already mentioned. Concerning other gauge fields, it seems to be appropriate to assign $F = SU(3)/U(2)$ to QCD gauge fields, $F = SL(2, C)/GL(1, C)$ to QED gauge fields and $F = SO(5)/SO(4)$ to weak interacting gauge fields. Some discussions concerned are following. In general, matter fields couple to $\mathbf{g}'$ -valued gauge fields. As for QCD, matter fields couple to the gauge fields of $U(2)$ subgroup but $SU(3)$ contains, as is well known, three types of $SU(2)$ subgroups and then after all they couple to all members of $SU(3)$ gauge fields. In case of QED, $GL(1, C)$ is locally isomorphic with $C^1 \cong U(1) \otimes R$. Then usual Abelian gauge fields are assigned to $U(1)$ subgroup of $GL(1, C)$. Georgi and Glashow suggested that the reason why the electric charge is quantized comes from the fact that $U(1)$ electromagnetic gauge group is a unfactorized subgroup of $SU(5)$ [9]. Our model is in the same situation because $GL(1, C)$ a unfactorized subgroup of $SL(2, C)$. For usual electromagnetic $U(1)$ gauge group, the electric charge unit “ e ” ($e > 0$) is for *one generator* of $U(1)$ but in case of $SL(2, C)$ which has *six generators*, the minimal unit of electric charge shared per one generator must be “ $e/6$ ”. This suggests that quarks and leptons might have the substructure simply because e , $2e/3$, $e/3 > e/6$. Finally as for weak interactions we adopt $F = SO(5)/SO(4)$.

It is well known that $SO(4)$ is locally isomorphic with $SU(2) \otimes SU(2)$. Therefore it is reasonable to think it the left-right symmetric gauge group : $SU(2)_L \otimes SU(2)_R$. As two $SU(2)$ s are direct product, it is able to have coupling constants $(\mathbf{g}_L, \mathbf{g}_R)$ independently. This is convenient to explain the fact of the disappearance of right-handed weak interactions in the low-energy region. Possibility of composite structure of quarks and leptons suggested by above $SL(2, C)$ -QED would introduce the thought that the usual left-handed weak interactions are intermediated by massive composite vector bosons as ρ -meson in QCD and that they are residual interactions due to substructure dynamics of quarks and leptons. The elementary massless gauge fields relate essentially to the structure of the real space-time manifold as the connection fields but on the other hand the composite vector bosons have nothing to do with it. Considering these discussions, we shall set the assumption : *“All kinds of gauge fields are elementary massless fields, belonging to spontaneously unbroken $SU(3)_C \otimes SU(2)_L \otimes SU(2)_R \otimes U(1)_{e.m}$ gauge group and quarks and leptons and $\mathbf{W}, \mathbf{Z}$ are all composite objects of the elementary matter fields.”*

3 Composite model

Our direct motivation towards compositeness of quarks and leptons is one of the results of the arguments in Sect.2, that is, $e, 2e/3, e/3 > e/6$. However, other several phenomenological facts tempt us to consider a composite model, e.g., repetition of generations, quark-lepton parallelism of weak isospin doublet structure, quark-flavor-mixings, etc. . Especially Bjorken[10]’s and Hung and Sakurai[11]’s suggestion of an alternative to unified weak-electromagnetic gauge theories have invoked many studies of composite models including composite weak bosons[12-17]. Our model is in the line of those studies. There are two ways to make composite models, that is, “Preons are all fermions.” or “Preons are both fermions and bosons (FB-model).” The merit of the former is that it can avoid the problem of a quadratically divergent self-mass of elementary scalar fields. However, even in the latter case such a disease is overcome if both fermions and bosons are the supersymmetric pairs, both of which carry the same quantum numbers except for the nature of Lorentz transformation (spin-1/2 or spin-0)[18]. Pati and Salam have suggested that the construction of a neutral composite object (neutrino in practice) needs both kinds of preons, fermionic as well as bosonic, if they carry the same charge for the Abelian gauge or belong to the same (fundamental) representation for the non-Abelian gauge[19]. This is a very attractive idea for

constructing the minimal model. Further, from the representation theory of Poincaré group both integer and half-integer spin angular momentum occur equally for massless particles[20]. If nature chooses “fermionic monism”, there must exist the additional special reason to select it. Then in this point also, the thought of the FB-model is minimal. Based on such considerations we proposed a FB-model of “*only one kind of spin-1/2 elementary field (Λ) and of spin-0 elementary field (Θ)*”[5]. Both have the same electric charge of “ $e/6$ ” (Maki has first proposed the FB-model with the universal electric charge $e/6$. [21])¹ and the same transformation properties of the fundamental representation (3, 2, 2) under the spontaneously unbroken gauge symmetry of $SU(3)_C \otimes SU(2)_L \otimes SU(2)_R$ (let us call $SU(2)_L \otimes SU(2)_R$ “hypercolor gauge symmetry”). Then Λ and Θ come into the supersymmetric pair which guarantees ’tHooft’s naturalness condition[22]. The $SU(3)_C$, $SU(2)_L$ and $SU(2)_R$ gauge fields cause the confining forces with confining energy scales of $\Lambda_c \ll \Lambda_L < (or \cong) \Lambda_R$ (Schrempp and Schrempp discussed them elaborately in Ref.[17]). Here we call positive-charged primons (Λ , Θ) “*matter*” and negative-charged primons ($\bar{\Lambda}$, $\bar{\Theta}$) “*antimatter*”. Our final goal is to build quarks, leptons and $\mathbf{W}, \mathbf{Z}$ from primons : Λ ($\bar{\Lambda}$) and Θ ($\bar{\Theta}$). Let us discuss that scenario next.

At the very early stage of the development of the universe, the matter fields (Λ , Θ) and their antimatter fields ($\bar{\Lambda}$, $\bar{\Theta}$) must have broken out from the vacuum. After that they would have combined with each other as the universe was expanding. That would be the first step of the existence of composite matters. There are ten types of them :

<i>spin</i> 1/2	<i>spin</i> 0	<i>e.m.charge</i>	<i>Y.M.representation</i>
$\Lambda\Theta$	$\Lambda\Lambda, \Theta\Theta$	$e/3$	$(\bar{3}, 1, 1) (\bar{3}, 3, 1) (\bar{3}, 1, 3), (7a)$
$\Lambda\bar{\Theta}, \bar{\Lambda}\Theta$	$\Lambda\bar{\Lambda}, \Theta\bar{\Theta}$	0	$(1, 1, 1) (1, 3, 1) (1, 1, 3), (7b)$
$\bar{\Lambda}\bar{\Theta}$	$\bar{\Lambda}\bar{\Lambda}, \bar{\Theta}\bar{\Theta}$	$-e/3$	$(3, 1, 1) (3, 3, 1) (3, 1, 3) .(7c)$

In this step the confining forces are, in kind, in $SU(3) \otimes SU(2)_L \otimes SU(2)_R$ gauge symmetry but the $SU(2)_L \otimes SU(2)_R$ confining forces must be main because of the energy scale of $\Lambda_L, \Lambda_R \gg \Lambda_c$ and then the color gauge coupling α_s and e.m. coupling constant α are negligible. As is well known, the coupling constant of $SU(2)$ confining force are characterized by $\varepsilon_i = \sum_a \sigma_\alpha^a \sigma_\beta^a$, where σ s are 2×2 matrices of $SU(2)$, $a = 1, 2, 3$, $\alpha, \beta = \Lambda, \bar{\Lambda}, \Theta, \bar{\Theta}$, $i = 0$ for singlet and $i = 3$ for triplet. They are calculated as $\varepsilon_0 = -3/4$ which causes the attractive force and $\varepsilon_3 = 1/4$ causing the repulsive

¹The notations of Λ and Θ are inherited from those in Ref.[21]. After this we shall call Λ and Θ “**Primon**” named by Maki which means “primordial particle”.

force. As concerns, $SU(3)_C$ octet and sextet states are repulsive but singlet, triplet and antitriplet states are attractive and then the formers are disregarded. Like this, two primons are confined into composite objects with more than one singlet state of any $SU(3)_C \otimes SU(2)_L \otimes SU(2)_R$. Note that three primon systems cannot make the singlet states of $SU(2)$. Then we omit them. In Eq.(7,b), the $(1, 1, 1)$ -state is the “most attractive channel”. Therefore $(\Lambda\bar{\Theta})$, $(\bar{\Lambda}\Theta)$, $(\Lambda\bar{\Lambda})$ and $(\Theta\bar{\Theta})$ of $(1, 1, 1)$ -states with neutral e.m. charge must have been most abundant in the universe. Further $(\bar{3}, 1, 1)$ - and $(3, 1, 1)$ -states in Eq.(7,a,c) are next attractive. They presumably go into $\{(\Lambda\Theta)(\bar{\Lambda}\bar{\Theta})\}$, $\{(\Lambda\Lambda)(\bar{\Lambda}\bar{\Lambda})\}$, etc. of $(1, 1, 1)$ -states with neutral e.m. charge. These objects may be the candidates for the “cold dark matters” if they have even tiny masses. It is presumable that the ratio of the quantities between the ordinary matters and the dark matters firstly depends on the color and hypercolor charges (maybe the ratio is more than $1/3 \times 3$). Finally the $(*, 3, 1)$ - and $(*, 1, 3)$ -states are remained ($*$ is $1, 3, \bar{3}$). They are also stable because $|\varepsilon_0| > |\varepsilon_3|$. They are, so to say, the “intermediate clusters” towards constructing ordinary matters,² namely quarks, leptons and $\mathbf{W}, \mathbf{Z}$. Here we call such intermediate clusters “subquarks” and denote them as follows :

	<i>Y.M.representation</i>	<i>spin</i>	<i>e.m.charge</i>	
$\alpha = (\Lambda\Theta),$	$\alpha_L : (\bar{3}, 3, 1), \quad \alpha_R : (\bar{3}, 1, 3)$	1/2	$e/3,$	(8a)
$\beta = (\Lambda\bar{\Theta}),$	$\beta_L : (1, 3, 1), \quad \beta_R : (1, 1, 3)$	1/2	0,	(8b)
$\mathbf{x} = (\Lambda\Lambda, \Theta\Theta),$	$\mathbf{x}_L : (\bar{3}, 3, 1), \quad \mathbf{x}_R : (\bar{3}, 1, 3)$	0	$e/3,$	(8c)
$\mathbf{y} = (\Lambda\bar{\Lambda}, \Theta\bar{\Theta}),$	$\mathbf{y}_L : (1, 3, 1), \quad \mathbf{y}_R : (1, 1, 3)$	0	0,	(8d)

and there are also their antisubquarks[16].³

Now we come to the step to build quarks and leptons. The gauge symmetry of the confining forces in this step is also $SU(2)_L \otimes SU(2)_R$ because the subquarks are in the triplet states of $SU(2)_{L,R}$ and then they are combined into singlet states by the decomposition of $3 \times 3 = 1 + 3 + 5$ in $SU(2)$. We make the first generation as follows :

	<i>e.m.charge</i>	<i>Y.M.representation</i>	
$< \mathbf{u}_i = < \alpha_i \mathbf{x}_i $	$2e/3$	$(3, 1, 1),$	(9a)
$< \mathbf{d}_i = < \bar{\alpha}_i \bar{\mathbf{x}}_i \mathbf{x}_i $	$-e/3$	$(3, 1, 1),$	(9b)
$< \nu_i = < \alpha_i \bar{\mathbf{x}}_i $	0	$(1, 1, 1),$	(9c)

²Such thoughts have been proposed by Maki in Ref.[21]

³The notations of $\alpha, \beta, \mathbf{x}$ and $\mathbf{y}$ are inherited from those in Ref.[16] written by Fritzsch and Mandelbaum, because ours is, in the subquark level, similar to theirs with two fermions and two bosons.

$$|< \mathbf{e}_i| = |< \bar{\alpha}_i \bar{\mathbf{x}}_i \bar{\mathbf{y}}_i| - e \quad (1, 1, 1), \quad (9d)$$

where i stands for L or R [12].⁴ Here we note that β and $\mathbf{y}$ do not appear. In practice $((\beta\mathbf{y}) : (1, 1, 1))$ -particle is a candidate for neutrino. But as Bjorken has pointed out[10], non-vanishing charge radius of neutrino is necessary for obtaining the correct low-energy effective weak interaction Lagrangian[17]. Therefore β is assumed not to contribute to forming quarks and leptons. Presumably composite $(\beta\beta)$ -; $(\beta\bar{\beta})$ -; $(\bar{\beta}\bar{\beta})$ -states may go into the dark matters. It is also noticeable that in this model the leptons have finite color charge radius and then $SU(3)$ gluons interact directly with the leptons at energies of the order of, or larger than Λ_L or Λ_R [18]. Concerning the confinements of primons and subquarks, the confining forces of two steps are in the same spontaneously unbroken $SU(2)_L \otimes SU(2)_R$ gauge symmetry. Here let us assume that subquarks in quarks are confined at the energy of 1.6 TeV (if admitting CDF's data[23]).

Concerning the running coupling constant of the $SU(2)$ gauge theory ($\alpha_W(Q^2)$) we know the following equation :

$$1/\alpha_W(Q^2) = b_2 \ln(Q/\Lambda_q)^2, \quad (10a)$$

$$b_2 = 1/(4\pi)\{22/3 - (2/3)N_f - (1/12)N_s\}, \quad (10b)$$

where Q is the effective energy of $SU(2)$ gluon exchange, Λ_q is the confinement scale of subquarks inside quarks and $N_f(N_s)$ is the numbers of fermions (scalars) contributing to the vacuum polarizations. We calculate $b_2 = 0.35$ which comes from that the number of confined fermionic subquarks are 4 ($\alpha_i, i = 1, 2, 3$ for color freedom, β) and 4 for bosons ($\mathbf{x}_i, \mathbf{y}$) contributing to the vacuum polarization. Using $b_2 = 0.35$ we get $\alpha_W = 0.040$ at $Q=10^{19}$ GeV and extrapolating from this value we obtain the result that the confining energy of primons (Λ, Θ) is 1.6×10^2 TeV, where we use $b_2 = 0.41$ which is calculated with three kinds of Λ and Θ owing to three color freedoms. In sum, the radii of $\alpha, \beta, \mathbf{x}$ and $\mathbf{y}$ are the inverse of 1.6×10^2 TeV and the radii of quarks are the inverse of 1.6 TeV.

Next let us see the higher generations. Harari and Seiberg have stated that the orbital and radial excitations seem to have the wrong energy scale (order of $\Lambda_{L,R}$) and then the most likely type of excitations is the addition of preon-antipreon pairs[13,24].

⁴Subquark configurations in Eq.(9) are essentially the same as those in Ref.[12] written by Królikowski, who proposed the model of one fermion and one boson with the same charge $e/3$.

Then using $\mathbf{y}_{L,R}$ in Eq.(8,d) we construct them as follows :

$$\begin{cases} \langle \mathbf{c} | &= \langle \alpha \mathbf{x} \mathbf{y} | \\ \langle \mathbf{s} | &= \langle \bar{\alpha} \mathbf{x} \mathbf{x} \mathbf{y} |, \end{cases} \quad \begin{cases} \langle \nu_\mu | &= \langle \alpha \bar{\mathbf{x}} \mathbf{y} | \\ \langle \mu | &= \langle \bar{\alpha} \mathbf{x} \mathbf{x} \mathbf{y} |, \end{cases} \quad \text{2nd generation (11a)}$$

$$\begin{cases} \langle \mathbf{t} | &= \langle \alpha \mathbf{x} \mathbf{y} \mathbf{y} | \\ \langle \mathbf{b} | &= \langle \bar{\alpha} \mathbf{x} \mathbf{x} \mathbf{y} \mathbf{y} |, \end{cases} \quad \begin{cases} \langle \nu_\tau | &= \langle \bar{\alpha} \bar{\mathbf{x}} \mathbf{y} \mathbf{y} | \\ \langle \tau | &= \langle \bar{\alpha} \mathbf{x} \mathbf{x} \mathbf{y} \mathbf{y} |, \end{cases} \quad \text{3rd generation, (11b)}$$

where the suffix L, R s are omitted for brevity. We can also make vector and scalar particles with (1,1,1) :

$$\begin{cases} \langle \mathbf{W}^+ | &= \langle \alpha^\dagger \alpha^\dagger \mathbf{x} | \\ \langle \mathbf{W}^- | &= \langle \bar{\alpha}^\dagger \bar{\alpha}^\dagger \bar{\mathbf{x}} |, \end{cases} \quad \begin{cases} \langle \mathbf{Z}_1^0 | &= \langle \alpha^\dagger \bar{\alpha}^\dagger | \\ \langle \mathbf{Z}_2^0 | &= \langle \alpha^\dagger \bar{\alpha}^\dagger \mathbf{x} \bar{\mathbf{x}} |, \end{cases} \quad \text{Vector (12a)}$$

$$\begin{cases} \langle \mathbf{S}^+ | &= \langle \alpha^\dagger \alpha^\dagger \mathbf{x} | \\ \langle \mathbf{S}^- | &= \langle \bar{\alpha}^\dagger \bar{\alpha}^\dagger \mathbf{x} |, \end{cases} \quad \begin{cases} \langle \mathbf{S}_1^0 | &= \langle \alpha^\dagger \bar{\alpha}^\dagger | \\ \langle \mathbf{S}_2^0 | &= \langle \alpha^\dagger \bar{\alpha}^\dagger \mathbf{x} \bar{\mathbf{x}} |, \end{cases} \quad \text{Scalar, (12b)}$$

where the suffix L, R s are omitted for brevity and $\uparrow, \downarrow$ indicate *spin up, spin down* states. They play the role of intermediate bosons same as π, ρ in the strong interactions. As Eq.(9) and Eq.(12) contain only α and $\mathbf{x}$ subquarks, we can draw the “*line diagram*”s of weak interactions as seen in Fig (1). Eq.(9,d) shows that the electron is constructed from antimatters only. We know, phenomenologically, that this universe is mainly made of protons, electrons, neutrinos, antineutrinos and unknown dark matters. It is said that protons and electrons in the universe are almost same in quantity. Our model show that one proton has the configuration of $(\mathbf{uud}) = (2\alpha, \bar{\alpha}, 3\mathbf{x}, \bar{\mathbf{x}})$; electron : $(\bar{\alpha}, 2\bar{\mathbf{x}})$; neutrino : $(\alpha, \bar{\mathbf{x}})$; antineutrino : $(\bar{\alpha}, \mathbf{x})$ and the dark matters are presumably constructed from the same amount of matters and antimatters because of their neutral charges. Therefore these facts may lead the thought that “the universe is the matter-antimatter-even object.” And then there exists a conception-leap between “proton-electron abundance” and “matter abundance” if our composite scenario is admitted (as for the possible way to realize the proton-electron excess universe, see Ref.[5]).

Our composite model contains two steps, namely the first is “subquarks made of primons” and the second is “quarks and leptons made of subquarks”. Here let us discuss about the mass generation mechanism of quarks and leptons as composite objects. Our model has only one kind of fermion : Λ and boson : Θ . The first step of “subquarks made of primons” seems to have nothing to do with ’tHooft’s anomaly matching condition[22] because there is no global symmetry with Λ and Θ . Therefore from this line of thought it is impossible to say anything about that $\alpha, \beta, \mathbf{x}$ and $\mathbf{y}$ are massless or massive. However, if it is the case that the neutral (1,1,1)-states

of primon-antiprion composites (as is stated above) become the dark matters, the masses of them presumably be less than the order of MeV from the phenomenological aspects of astrophysics. Then we may assume that these subquarks are massless or almost massless compared with $\Lambda_{L,R}$ in practice, that is, utmost a few MeV. In the second step, the arguments of 'tHooft's anomaly matching condition are meaningful. The confining of subquarks must occur at the energy scale of $\Lambda_{L,R} \gg \Lambda_c$ and then it is natural that $\alpha_s, \alpha \rightarrow 0$ and that the gauge symmetry group is the spontaneously unbroken $SU(2)_L \otimes SU(2)_R$ gauge group. Seeing Eq.(9), we find quarks and leptons are composed of the mixtures of subquarks and antishquarks. Therefore it is proper to regard subquarks and antishquarks as different kinds of particles. From Eq.(8,a,b) we find eight kinds of fermionic subquarks (3 for $\alpha, \bar{\alpha}$ and 1 for $\beta, \bar{\beta}$). So the global symmetry concerned is $SU(8)_L \otimes SU(8)_R$. Then we arrange :

$$(\beta, \bar{\beta}, \alpha_i, \bar{\alpha}_i \quad i = 1, 2, 3)_{L,R} \quad \text{in} \quad (SU(8)_L \otimes SU(8)_R)_{global}, \quad (13)$$

where i s are color freedoms. Next, the fermions in Eq.(12) are confined into the singlet states of the local $SU(2)_L \otimes SU(2)_R$ gauge symmetry and make up quarks and leptons as seen in Eq.(9) (eight fermions). Then we arrange :

$$(\nu_{\mathbf{e}}, \mathbf{e}, \mathbf{u}_i, \mathbf{d}_i \quad i = 1, 2, 3)_{L,R} \quad \text{in} \quad (SU(8)_L \otimes SU(8)_R)_{global}, \quad (14)$$

where i s are color freedoms. From Eq.(13) and Eq.(14) the anomalies of the subquark level and the quark-lepton level are matched and then all composite quarks and leptons (in the 1st generation) are remained massless. Note again that presumably, β and $\bar{\beta}$ in Eq.(12) are composed into "bosonic" $(\beta\beta)$, $(\beta\bar{\beta})$ and $(\bar{\beta}\bar{\beta})$, which vapor out to the dark matters. Schrempp and Schrempp have discussed about a confining $SU(2)_L \otimes SU(2)_R$ gauge model with three fermionic preons and stated that it is possible that not only the left-handed quarks and leptons are composite but also the right-handed are so on the condition that Λ_R/Λ_L is at least of the order of 3[17]. If CDF's data[23] truly indicates the compositeness of quarks, Λ_L is presumably around 1.6 TeV. As seen in Eq.(12.a) the existence of composite $\mathbf{W}_R, \mathbf{Z}_R$ is predicted. As concerning, the fact that they are not observed yet means that the masses of $\mathbf{W}_R, \mathbf{Z}_R$ are larger than those of $\mathbf{W}_L, \mathbf{Z}_L$ and that $\Lambda_R > \Lambda_L$. Owing to 'tHooft's anomaly matching condition the small mass nature of the 1st generation comparing to Λ_L is guaranteed but the evidence that the quark masses of the 2nd and the 3rd generations become larger as the generation numbers increase seems to have nothing to do with the anomaly matching mechanism in our model, because as seen in Eq.(11,a,b) these generations are obtained by just adding

scalar $\mathbf{y}$ -particles. This is different from Abott and Farhi's model in which all fermions of three generations are equally embedded in $SU(12)$ global symmetry group and all members take part in the anomaly matching mechanism[15,25]. Concerning this, let us discuss a little about subquark dynamics inside quarks. According to "Uncertainty Principle" the radius of the composite particle is, in general, roughly inverse proportional to the kinetic energy of the constituent particles moving inside it. The radii of quarks may be around $1/\Lambda_{L,R}$. So the kinetic energies of subquarks may be more than hundreds GeV and then it is considered that the masses of quarks essentially depend on the kinetic energies of subquarks and such a large binding energy as counterbalances them. As seen in Eq.(10,a,b) our model shows that the more the generation number increases the more the number of the constituent particles increases. So assuming that the radii of all quarks do not vary so much (because we have no experimental evidences yet), the interaction length among subquarks inside quarks becomes shorter as generation numbers increase and accordingly the average kinetic energy per one subquark may increase. Therefore integrating out the details of subquark dynamics it could be said that the essential feature of increasing masses of the 2nd and the 3rd generations is simply because their masses are described as a increasing function of the sum of the kinetic energies of constituent subquarks. From the Review of Particles and Fields[29] we can phenomenologically parameterized the mass spectrum of quarks and leptons as follows :

$$M_{UQ} = 1.2 \times 10^{-4} \times (10^{2.05})^n \quad \text{GeV} \quad \text{for } \mathbf{u, c, t}, \quad (15a)$$

$$M_{DQ} = 3.0 \times 10^{-4} \times (10^{1.39})^n \quad \text{GeV} \quad \text{for } \mathbf{d, s, b}, \quad (15b)$$

$$M_{DL} = 3.6 \times 10^{-4} \times (10^{1.23})^n \quad \text{GeV} \quad \text{for } \mathbf{e, \mu, \tau}, \quad (15c)$$

where $n = 1, 2, 3$ are the generation numbers. They seem to be geometric-ratio-like. The slopes of the up-quark sector and down-quark sector are different, so it seems that each has different aspects in subquark dynamics. From Eq.(15) we obtain $M_{\mathbf{u}} = 13.6$ MeV, $M_{\mathbf{d}} = 7.36$ MeV and $M_{\mathbf{e}} = 6.15$ MeV. These are a little unrealistic compared with the experiments[4]. But considering the above discussions about the anomaly matching conditions (Eq.(13,14)), it is natural that the masses of the members of the 1st generation are roughly equal to those of the subquarks, that is, a few MeV. The details of their mass-values depend on the subquark dynamics owing to the effects of electromagnetic and color gauge interactions. These mechanism has studied by Weinberg[26] and Fritzsch[27].

One of the experimental evidences inspiring the SM is the “universality” of the coupling strength among the weak interactions. Of course if the intermediate bosons are gauge fields, they couple to the matter fields universally. But the inverse of this statement is not always true, namely the quantitative equality of the coupling strength of the interactions does not necessarily imply that the intermediate bosons are elementary gauge bosons. In practice the interactions of ρ and ω are regarded as indirect manifestations of QCD. In case of chiral $SU(2) \otimes SU(2)$ the pole dominance works very well and the predictions of current algebra and PCAC seem to be fulfilled within about 5%[18]. Fritzsche and Mandelbaum[16,18] and Gounaris, K ogerler and Schildknecht[28,29] have elaborately discussed about universality of weak interactions appearing as a consequence of current algebra and **W**-pole dominance of the weak spectral functions from the stand point of the composite model. Extracting the essential points from their arguments we shall mention the followings .

In the first generation let the weak charged currents be written in terms of the subquark fields as :

$$\mathbf{J}_\mu^+ = \overline{U} h_\mu D, \quad \mathbf{J}_\mu^- = \overline{D} h_\mu U, \quad (16)$$

where $U = (\alpha \mathbf{x})$, $D = (\overline{\alpha} \mathbf{x})$ and $h_\mu = \gamma_\mu(1 - \gamma_5)$. Reasonableness of Eq.(16) may given by the fact that $M_W \ll \Lambda_{L,R}$ (where M_W is **W**-boson mass). Further, let U and D belong to the doublet of the global weak isospin $SU(2)$ group and $\mathbf{W}^+$, $\mathbf{W}^-$, $1/\sqrt{2}(\mathbf{Z}_1^0 - \mathbf{Z}_2^0)$ be in the triplet and $1/\sqrt{2}(\mathbf{Z}_1^0 + \mathbf{Z}_2^0)$ be in the singlet of $SU(2)$. These descriptions seem to be natural if we refer the diagrams in Fig.(1). The universality of the weak interactions are inherited from the universal coupling strength of the algebra of the global weak isospin $SU(2)$ group with the assumption of **W**-, **Z**-pole dominance. The universality including the 2nd and the 3rd generations are based on the above assumptions and the concept of the flavor-mixings. The quark-flavor-mixings in the weak interactions are expressed by Cabbibo-Kobayashi-Maskawa (CKM)-matrix (V_{ij}) based on the SM. Its nine matrix elements (in case of three generations) are free parameters (in practice four parameters with the unitarity) and this point is said to be one of the drawback of the SM along with non-understanding of the origins of the quark-lepton mass spectrum and generations. In the SM, the quark fields (lepton fields also) are elementary and then we are able to investigate, at the utmost, the external relationship among them. On the other hand if quarks are the composites of substructure constituents, the quark-flavor-mixing phenomena must be understood by the substructure dynamics and the values of CKM matrix elements become materials

for studying these . In our model “*the quark-flavor-mixings occur by creations or annihilations of $\mathbf{y}$ -particles inside quarks*”. The $\mathbf{y}$ -particle is a neutral scalar subquark in the $\mathbf{3}$ -state of $SU(2)_L$ group and then couples to two hypercolor gluons (denoted by $\mathbf{g}_h$). By this mechanism we obtained $|V_{ub}| = 3.45 \times 10^{-3}$, $|V_{ts}| = 2.62 \times 10^{-2}$ and $|V_{td}| = 1.40 \times 10^{-3}$ (for detailed analysis see Ref.[52]).

4 Mass differences and CP-Violations in Neutral Meson Systems

a. Mass difference ΔM_P by $P^0 - \overline{P}^0$ mixing

The typical $\Delta F = 2$ phenomenon is the mixing between a neutral pseudo scalar meson (P^0) and its antimeson ($\overline{P}^0$). There are six types of them, e.g., $K^0 - \overline{K}^0$, $D^0 - \overline{D}^0$, $B_d^0 - \overline{B}_d^0$, $B_s^0 - \overline{B}_s^0$, $T_u^0 - \overline{T}_u^0$ and $T_c^0 - \overline{T}_c^0$ mixings. Usually they have been considered to be the most sensitive probes of higher-order effects of the weak interactions in the SM. The basic tool to investigate them is the “box diagram”. By using this diagram to the K_L - K_S mass difference, Gaillard and Lee predicted the mass of the charm quark[30]. Later, Wolfenstein suggested that the contribution of the box diagram which is called the short-distance (SD) contribution cannot supply the whole of the mass difference ΔM_K and there are significant contributions arising from the long-distance (LD) contributions associated with low-energy intermediate hadronic states[31]. As concerns, the LD-phenomena occur in the energy range of few hundred MeV and the SD-phenomena around 100 GeV region. Historically there are various investigations for P^0 - $\overline{P}^0$ mixing problems[3][32-41] and many authors have examined them by use of LD- and SD-contributions. In summary, the comparison between the theoretical results and the experiments about ΔM_P ($P = K, D$ and B_d) are as follows :

$$\Delta M_K^{LD} \approx \Delta M_K^{SD} \approx \Delta M_K^{exp}, \quad (17a)$$

$$\Delta M_D^{SD} \ll \Delta M_D^{LD} (\ll \Delta M_D^{exp}, upper\ bound), \quad (17b)$$

$$\Delta M_{B_d}^{LD} \ll \Delta M_{B_d}^{SD} \simeq \Delta M_{B_d}^{exp}. \quad (17c)$$

Concerning Eq.(17a) it is explain that $\Delta M_K = \Delta M_K^{SD} + D\Delta M_K^{LD}$ where “ D ” is a numerical value of order $O(1)$. As for Eq(17c), they found that $\Delta M_{B_d}^{LD} \approx 10^{-16}$ GeV

and $\Delta M_{B_d}^{SD} \approx 10^{-13}$ GeV, then the box diagram is the most important for $B_d^0\text{-}\overline{B}_d^0$ mixing. Computations of $\Delta M_{B_d}^{SD}$ and $\Delta M_{B_s}^{SD}$ from the box diagrams in the SM give

$$\Delta M_{B_s}^{SD}/\Delta M_{B_d}^{SD} \simeq (M_{B_s}/M_{B_d})|V_{ts}/V_{td}|^2(B_{B_s}f_{B_s}^2/B_{B_d}f_{B_d}^2)\zeta, \quad (18)$$

where V_{ij} s stand for CKM matrix elements; M_P : P-meson mass; ζ : a QCD correction of order $O(1)$; B_B : Bag factor of B-meson and f_B : decay constant of B-meson. Measurements of $\Delta M_{B_d}^{exp}$ and $\Delta M_{B_s}^{exp}$ are, therefore, said to be useful to determine $|V_{ts}/V_{td}|$ [42][43]. Concerning Eq.(17b), they found that $\Delta M_D^{LD} \approx 10^{-15}$ GeV and $\Delta M_D^{SD} \approx 10^{-17}$ GeV[3][37] but the experimental measurement is $\Delta M_D^{exp} < 1.3 \times 10^{-13}$ GeV[4]. Further there is also a study that ΔM_D^{LD} is smaller than 10^{-15} GeV by using the heavy quark effective theory[38]. Then many people state that it would be a signal of new physics beyond the SM if the future experiments confirm that $\Delta M_D^{exp} \simeq 10^{-14} \sim 10^{-13}$ GeV[32-38]. Above investigations are based on the calculations of SD-contributions with (or without) LD-contributions in the SM.

On the other hand some authors have studied these phenomena in the context of the theory explained by the single dynamical origin. Cheng and Sher[44], Liu and Wolfenstein[40], and Gérard and Nakada[41] have thought that all $P^0\text{-}\overline{P}^0$ mixings occur only by the dynamics of the TeV energy region which is essentially the same as the super-weak (SW) idea originated by Wolfenstein[2]. They extended the original SW-theory (which explains CP violation in the K -meson system) to other flavors by setting the assumption that $\Delta F = 2$ changing neutral *spin* 0 particle with a few TeV mass (denoted by H) contributes to the “real part” of M_{ij} which determines ΔM_P and also the “imaginary part” of M_{ij} which causes the indirect CP violation. The ways of extensions are that H -particles couple to quarks by the coupling proportional to $\sqrt{m_i m_j}$ [40][44], $(m_i/m_j)^n$ $n = 0, 1, 2$ [40] and $(m_i + m_j)$ [41] where i, j are flavors of quarks coupling to H . It is suggestive that the SW-couplings depend on quark masses (this idea is adopted in our model discussed below). Cheng and Sher[44] and Liu and Wolfenstein[40] obtained that $\Delta M_D = (m_c/m_s)\Delta M_K^{exp} \approx 10^{-14}$ GeV with the assumption that H -exchange mechanism saturates the ΔM_K^{exp} bound, which is comparable to $\Delta M_D^{exp} < 1.3 \times 10^{-13}$ GeV[4]. Concerning B -meson systems they found that $\Delta M_{B_s}/\Delta M_{B_d} = m_s/m_d \simeq 20$ which seems agreeable to $(\Delta M_{B_s}/\Delta M_{B_d})_{exp} > 13$ [43]. However using their scheme it is calculated that

$$\Delta M_{B_d}/\Delta M_K = (B_{B_d}f_{B_d}^2/B_Kf_K^2)(M_{B_d}/M_K)(m_b/m_s) \simeq 300, \quad (19)$$

where we use $m_b = 4.3$ GeV, $m_s = 0.2$ GeV, $M_{B_d} = 5.279$ GeV, $M_K = 0.498$ GeV,

$B_{B_d}f_{B_d}^2 = (0.22\text{GeV})^2$, $B_Kf_K^2 = (0.17\text{GeV})^2$. It seems larger than $(\Delta M_{B_d}/\Delta M_K)_{exp} = 85.8 \pm 3.6[43]$. This result is caused by rather large b-quark mass value.

Now let us discuss P^0 - $\overline{P}^0$ mixings by using our FB-model. The discussions start from the assumption that the mass mixing matrix $M_{ij}(P)$ ($i(j) = 1(2)$ denotes $P^0(\overline{P}^0)$) is saturated by the super-weak-type interactions causing a direct $\Delta F = 2$ transitions. We usually calculate ΔM_P as

$$M_{12}(P) = \langle \overline{P}^0 | \mathcal{H}_{SW}^{\Delta F=2} | P^0 \rangle, \quad (20a)$$

$$\Delta M_P = M_H - M_L \simeq 2|M_{12}(P)|, \quad (20b)$$

where we assume $ImM_{12} \ll ReM_{12}$ which is experimentally preferable[3][45], and $M_{H(L)}$ stands for heavier (lighter) P^0 -meson mass. Applying the vacuum-insertion calculation to the hadronic matrix element as $\langle \overline{P}^0 | [\overline{q}_i \gamma_\mu (1 - \gamma_5) q_j]^2 | P^0 \rangle \sim B_P f_P^2 M_P^2 [3]$ we get

$$M_{12}(P) = (1/12\pi^2) B_P f_P^2 M_P \mathcal{M}_P. \quad (21)$$

The details of $\mathcal{M}_P$ are model-dependent, e.g., the box diagram in the SM; the neutral *spin* 0 particle exchange in the SW-theory. In case of our FB-model, the diagrams contributing to $\mathcal{M}_P$ are seen in Fig.(2). In our model P^0 - $\overline{P}^0$ mixings occur due to “**y**-exchange” between two quarks inside the present P^0 -meson. This is a kind of the realizations of Wolfenstein’s SW-idea[2]. The schematic illustration is as follows : two particles (quarks) with radius order of $1/\Lambda_q$ (a few TeV^{-1}) are moving to and fro inside a sphere (meson) with radius order of GeV^{-1} . The **y**-exchange interactions would occur when two quarks inside P^0 -meson interact in contact with each other because **y**-particles are confined inside quarks. As seen in Fig.(2), the contributions of **y**-exchanges seem common among various P^0 -mesons. Upon this, setting the assumption : “*universality of the y-exchange interactions*”, we rewrite $\mathcal{M}_P$ as

$$\mathcal{M}_P = n_P \eta(P) \tilde{\mathcal{M}}_i(P), \quad (22)$$

where $n_P = 1$ for $P = K, D, B_d, T_u$; $n_P = 2$ for $P = B_s, T_c$, $i = 1$ for K, D, B_s, T_u ; $i = 2$ for B_d, T_c . Then the universality means explicitly that

$$\tilde{\mathcal{M}}_1(K) = \tilde{\mathcal{M}}_1(D) = \tilde{\mathcal{M}}_1(B_s) = \tilde{\mathcal{M}}_1(T_c), \quad (23a)$$

$$\tilde{\mathcal{M}}_2(B_d) = \tilde{\mathcal{M}}_2(T_u). \quad (23b)$$

The explanation of n_P is such that K and D have one **y**-particle and one **y**-particle exchanges; B_d and T_u have two **y**-particles and both of them exchange simultaneously,

so for them we set $n_P = 1$ and B_s and T_c have two $\mathbf{y}$ -particles but one of them exchanges, so they have $n_P = 2$ because the probability becomes double. The “ i ” means the number of exchanging $\mathbf{y}$ -particles in the present diagram. Concerning $\eta(P)$, we shall explain as follows : In our FB-model P^0 - $\overline{P}^0$ mixing occurs by the “contact interaction” of two quarks coliding inside P^0 -meson. Therefore the probability of this interaction may be considered inverse proportional to the volume of the present P^0 -meson, e.g., the larger radius K -meson gains the less-valued probability of the coliding than the smaller radius D - (or B_s -) meson. The various aspects of hadron dynamics seem to be successfully illustrated by the semi-relativistic picture with “Breit-Fermi Hamiltonian”[46]. Assuming the power-law potential $V(r) \sim r^\nu$ (ν is a real number), the radius of P^0 -meson (denoted by $\mathbf{r}_P$) is proportional to $\mu_P^{-1/(2+\nu)}$, where μ_P is the reduced mass of two quark-masses inside P^0 -meson[46]. Then the volume of P^0 -meson is proportional to $\mathbf{r}_P^3 \sim \mu_P^{-3/(2+\nu)}$. After all we could assume for $\eta(P)$ in Eq.(22) as

$$\eta(P) = \xi(\mu_P/\mu_K)^1 \quad \text{for linear - potential,} \quad (24a)$$

$$= \xi(\mu_P/\mu_K)^{1.5} \quad \text{for log - potential,} \quad (24b)$$

where ξ is a dimensionless numerical factor depending on the details of the dynamics and $\eta(P)$ is normalized by μ_K (reduced mass of s - and d -quark in K meson) for convenience. We may think that the $\mathbf{y}$ -exchange is described by the overlapping of the wave functions of two quarks inside P^0 -meson. Then we shall write as

$$|\tilde{\mathcal{M}}_i(P)| = (1/\Lambda_q^2) |\kappa \int \Psi_q(r) \Psi_{q'}(r) d^3r|, \quad (25a)$$

$$\simeq |\kappa \int \Psi_q(r) \Psi_{q'}(r) d^3r| \times 10^{-7} \quad \text{GeV}^{-2}, \quad (25b)$$

where $\Psi_q(r)$ is a radial wave function of q -quark, κ is a dimensionless complex numerical factor caused by unknown subquark dynamics and may depend on $| < q' | \overline{\mathbf{y}}(\partial_\mu \mathbf{y}) | q > |$. In Eq.(25b) we estimate a few TeV as Λ_q .

From the experimental informations the complex $M_{12}(K)$ is evaluated as [3]

$$M_{12}^{exp}(K) = -(0.176 + i0.114 \times 10^{-2}) \times 10^{-14} \quad \text{Gev.} \quad (26)$$

On the other hand, setting $P=K$ in Eq.(21) we obtain

$$|\mathcal{M}_K| = |M_{12}(K)|/(1/12\pi^2) B_K f_K^2 M_K \simeq 0.15 \times 10^{-10} \quad \text{GeV}^{-2}, \quad (27)$$

where we use $|M_{12}(K)| = |M_{12}^{exp}(K)|$ from Eq.(26), $B_K f_K^2 = (0.17\text{GeV})^2$ and $M_K = 0.498 \text{ GeV}$. Further setting $P = K$ in Eq.(22), (24), and (25) we have

$$|\mathcal{M}_K| = \xi |\tilde{\mathcal{M}}_1(K)| = \xi |\kappa \int \Psi_s(r) \Psi_d(r) d^3r| \times 10^{-7} \quad \text{GeV}^{-2}, \quad (28)$$

From Eq.(27) and (28) we obtain

$$\xi|\kappa| \int \Psi_s(r)\Psi_d(r)d^3r| \simeq 10^{-4} \quad \text{GeV}^{-2}. \quad (29)$$

As it may be expected that

$$|\int \Psi_s(r)\Psi_d(r)d^3r| \simeq O(10^{-2}) \sim O(10^{-1}), \quad (30)$$

we have

$$\xi|\kappa| \simeq O(10^{-3}) \sim O(10^{-2}). \quad (31)$$

Eq.(30) has to be ascertained in future. If the above investigation is the case, our picture corresponds to the scheme of the Higgs (of a few TeV mass value) exchange in the SW-theory[40][44][47][49][50]. Note that the mass value of **y**-particle itself is less than a few MeV as seen in Sect.(3).

The experimental results of ΔM_P are as follows[4][43] :

$$\Delta M_K = (3.510 \pm 0.018) \times 10^{-15} \quad \text{GeV}, \quad (32a)$$

$$\Delta M_D < 1.3 \times 10^{-13} \quad \text{GeV}, \quad (32b)$$

$$\Delta M_{B_d} = (3.01 \pm 0.13) \times 10^{-13} \quad \text{GeV}, \quad (32c)$$

$$\Delta M_{B_s} > 4.0 \times 10^{-12} \quad \text{GeV}. \quad (32d)$$

Using Eq.(20), (21) and (32), we have

$$|\mathcal{M}_D| < 8.1|\mathcal{M}_K|, \quad (33a)$$

$$|\mathcal{M}_{B_d}| = 4.86|\mathcal{M}_K|, \quad (33b)$$

$$|\mathcal{M}_{B_s}| > 49.3|\mathcal{M}_K|. \quad (33c)$$

At the level of $\mathcal{M}_P$, it seems that

$$|\mathcal{M}_P|/|\mathcal{M}_K| \simeq O(1) \sim O(10), \quad (34)$$

where $P = D, B_d, B_s$.

Here let us go on to more precise investigations. In Eq.(23) let us extend the “universality” and assume that $|\tilde{\mathcal{M}}_1(K)| \simeq |\tilde{\mathcal{M}}_2(B_d)|$. Using Eq.(22), (24) and (33b) we obtain

$$\mu_{B_d}/\mu_K = 4.86 \quad \text{for linear - potential}, \quad (35a)$$

$$= 2.87 \quad \text{for log - potential}, \quad (35b)$$

where $B_{B_d}f_{B_d}^2 = (0.22\text{GeV})^2$, $B_Kf_K^2 = (0.17\text{GeV})^2$ are used. Note that, comparing with the case of Eq.(19), we can evade the large enhancement by b -quark mass effect. This is because the quark mass dependence is introduced through the reduced mass (in which the effect of heavier mass decreases). Some discussions are as follows : If we adopt the pure non-relativistic picture it may be that $\mu_K \simeq \mu_{B_d} \simeq m_d \simeq (\mu_D \simeq \mu_{T_u})$ but from the semi-relativistic standpoint it seems preferable that $\mu_K(< \mu_D) < \mu_{B_d}(< \mu_{T_u})$ because the effective mass value of “ d -quark” in B_d -meson is considered larger than that in K -meson. It may be caused by that the kinetic energy of “ d -quark” in B_d -meson is larger than that in K -meson owing to the fact : $\mathbf{r}_{B_d} < \mathbf{r}_K$. Then we can expect the plausibility of Eq.(35). Of course it may be also a question whether the extension of the universality : $|\tilde{\mathcal{M}}_1(K)| \simeq |\tilde{\mathcal{M}}_2(B_d)|$ is good or not (this point may influence Eq.(35)), which will become clear when the experimental result about ΔM_{T_u} is confirmed in future and compared with ΔM_{B_d} . Next, let us study ΔM_D . In order to estimate the lower limit of ΔM_D^{SW} (denoted by $(\Delta M_D^{SW})_{LL}$) we set $\mu_D = \mu_K$ tentatively in Eq.(24) and obtain

$$(\Delta M_D^{SW})_{LL} = 4.67 \times \Delta M_K = 1.6 \times 10^{-14} \quad \text{GeV}, \quad (36)$$

where we use $B_Df_D^2 = (0.19\text{GeV})^2$ and Eq.(20), (21), (22), (23a) and (32a). In the same way, assuming $\mu_D = 1.5 \times \mu_K$ for example and using Eq.(24) we have

$$\Delta M_D^{SW} = (2.9 \sim 5.4) \times 10^{-14} \quad \text{GeV}, \quad (37)$$

which is consistent and comparable with Eq.(32b). These values are similar to the results by Cheng and Sher[44] and Liu and Wolfenstein[40].

The study of ΔM_{B_s} is as follows. Both s - and b -quark in B_s -meson are rather massive and then supposing availability of the non-relativistic scheme we have

$$\mu_{B_s} = m_s m_b / (m_s + m_b) = 0.19 \quad \text{GeV}, \quad (38)$$

where $m_s = 0.2 \text{ GeV}$ and $m_b = 4.3 \text{ GeV}$ are used. If we adopt $\mu_K = 0.01 \text{ GeV} (\simeq m_d)$ for example we obtain

$$\eta(B_s) = 19.0\xi \quad \text{for linear - potential}, \quad (39a)$$

$$= 82.8\xi \quad \text{for log - potential}, \quad (39b)$$

By using Eq.(20b), (21), (22) and (23a) we have

$$\Delta M_{B_s}^{SW} = 2(B_{B_s}f_{B_s}^2 M_{B_s} \eta(B_s) / B_K f_K^2 M_K \eta(K)) \Delta M_K^{SW}, \quad (40)$$

where factor 2 comes from $n_{B_s} = 2$ in Eq.(22). Assuming that $\Delta M_K^{SW} = \Delta M_K^{exp}$ (that is, the super-weak exchange saturates the ΔM_K^{exp} bound) and using Eq.(39) we obtain

$$\Delta M_{B_s}^{SW} = (0.31 \sim 1.4) \times 10^{-11} \text{ GeV}, \quad (41)$$

where we use $B_{B_s} f_{B_s}^2 = (0.25 \text{ GeV})^2 [42]$. From Eq.(32c) and (41) we get

$$\Delta M_{B_s}^{SW} / \Delta M_{B_d}^{SW} = 10 \sim 50, \quad (42a)$$

$$x_s = \tau_{B_s} \Delta M_{B_s} = 8 \sim 30, \quad (42b)$$

where we set $\Delta M_{B_d}^{SW} = \Delta M_{B_d}^{exp}$ and use $\tau_{B_s} = 2.4 \times 10^{12} \text{ GeV}^{-1} [4]$. Note that the present experimental result is $\Delta M_{B_s}^{exp} / \Delta M_{B_d}^{exp} > 12 [43]$. If we adopt the box diagram calculation in the SM and use Eq.(18) with the unitary assumption of CKM-matrix elements, it is found that [42][43]

$$\Delta M_{B_s}^{SD} / \Delta M_{B_d}^{SD} = 10 \sim 100. \quad (43)$$

Therefore, from the studies of ΔM_{B_d} and ΔM_{B_s} it is difficult to clarify which scheme (SW or SD in the SM) is true, at least until the future experiments confirm the values of $|V_{ts}/V_{td}|$ and ΔM_{B_s} .

Finally let us estimate $\Delta M_{T_u}^{SW}$ and $\Delta M_{T_c}^{SW}$. Setting $\mu_{T_u} = \mu_D$ (though $\mu_{T_u} > \mu_D$ in practice) and using Eq.(20b), (21), (22), (23b) and (24) we estimate the lower limit of $\Delta M_{T_u}^{SW}$ (denoted by $(\Delta M_{T_u}^{SW})_{LL}$) as

$$(\Delta M_{T_u}^{SW})_{LL} = (B_{T_u} f_{T_u}^2 M_{T_u} / B_{B_d} f_{B_d}^2 M_{B_d}) \Delta M_{B_d}^{SW} = 7.2 \times 10^{-10} \text{ GeV}, \quad (44)$$

where we use $B_{T_u} f_{T_u}^2 = (1.9 \text{ GeV})^2 [3]$, $M_{B_d} = 5.279 \text{ GeV}$, $M_{T_u} = 170 \text{ GeV}$ and set $\Delta M_{B_d}^{SW} = \Delta M_{B_d}^{exp}$ in Eq.(31c). Note that $|\tilde{\mathcal{M}}_2(T_u)| = |\tilde{\mathcal{M}}_2(B_d)|$ is used in Eq.(44). Cheng and Sher's scheme [44] predicts $\Delta M_{T_u} \simeq 10^{-7} \text{ GeV}$ which is order 10^3 larger than Eq.(44). (In Ref.[44] they estimated $\Delta M_T \simeq 10^{-10} \text{ GeV}$ using rather small t -quark mass value). For evaluating ΔM_{T_c} , we calculate

$$\mu_{T_c} = m_c m_t / (m_c + t_b) = 1.34 \text{ GeV}, \quad (45)$$

where $m_c = 1.35 \text{ GeV}$ and $m_t = 170 \text{ GeV}$ are used. Then we get from Eq.(24)

$$\eta(T_c) = 134\xi \quad \text{for linear - potential}, \quad (46a)$$

$$= 1551\xi \quad \text{for log - potential}, \quad (46b)$$

where we set $\mu_K = 0.01$ GeV for example. After all with Eq.(46) we obtain

$$\Delta M_{T_c}^{SW} = 2(B_{T_c} f_{T_c}^2 M_{T_c} / B_K f_K^2 M_K) (\eta(T_c) / \eta(K)) \Delta M_K^{SW} = (4 \sim 47) \times 10^{-8} \text{ GeV}, \quad (47)$$

where we adopt $n_{T_c} = 2$, $B_{T_c} f_{T_c}^2 = (1.9 \text{ GeV})^2 [3]$, $M_{T_u} = 171.35$ GeV and $\Delta M_K^{SW} = \Delta M_K^{exp}$. Note that $|\tilde{\mathcal{M}}_1(T_c)| = |\tilde{\mathcal{M}}_1(K)|$ is used in Eq.(47).

b. CP violations in P^0 - $\overline{P}^0$ mixings

Presently observed CP violations are only in the Kaon system and are still not inconsistent with the SW-model though there exists the discrepancy between E731 and NA31 experiments concerning $Re(\epsilon'/\epsilon)_K$. If $Re(\epsilon'/\epsilon)_K = 0$ is confirmed by experiments, CKM-mixing matrix has no phase factor[48] and CP violation in K -systems can be explained only by the SW-theory. If $Re(\epsilon'/\epsilon)_K \neq 0$ is confirmed by experiments (which implies the existence of CP violation by decay modes), there are several standpoints, e.g., only the standard CKM-theory explains it with (or without) the LD-contributions; the SW-interactions coexist with the SM, etc.. On the other hand, at the present stage no evidence of CP violation in other heavier meson systems is found. As widely discussed, CP asymmetries in $B_s \rightarrow \psi K_s$ and $B \rightarrow 2\pi$ may give us the crucial clues[41][51].

In this paper we discuss CP violations by mass-mixings which are assumed to be saturated by the SW-interactions. In the CP conserving limit in the P^0 -meson systems, $M_{12}(P)$ s are supposed to be real positive. Note that $CP|P_H\rangle = -|P_H\rangle$ and $CP|P_L\rangle = |P_L\rangle$ where H (L) means heavy (light). If the CP violating SW-interactions are switched on, $M_{12}(P)$ becomes complex. Following Gérard and Nakada's notation[41][45], we write as

$$M_{12} = |M_{12}| \exp(i\theta_P), \quad (48)$$

with

$$\tan \theta_P = Im M_{12}(P) / Re M_{12}(P). \quad (49)$$

As we assume that the SW-interaction saturates CP violation, we can write

$$Im \langle \overline{P}^0 | H_{SW}^{\Delta F=2} | P \rangle = Im M_{12}(P). \quad (50)$$

From Eq.(20), (21) and (22) we obtain

$$Im M_{12}(P) = \mathcal{A} \cdot Im \tilde{\mathcal{M}}_i(P), \quad (51)$$

where $\mathcal{A} = (1/12\pi^2) B_P f_P^2 M_P \eta(P)$. Therefore the origin of CP violation of P^0 -meson system is only in $\tilde{\mathcal{M}}_i(P)$. The Factor “ $\mathcal{A}$ ” in Eq.(51) is common also in $Re M_{12}(P)$ and then we have

$$Im M_{12}(P) / Re M_{12}(P) = Im \tilde{\mathcal{M}}_i(P) / Re \tilde{\mathcal{M}}_i(P). \quad (52)$$

If the universality of Eq.(23) is admitted, we obtain

$$\theta_K = \theta_D = \theta_{B_s} = \theta_{T_c}, \quad (53a)$$

$$\theta_{B_d} = \theta_{T_u}, \quad (53b)$$

These are the predictions about CP violations from the stand point of our FB-model. Whether θ_K and θ_{B_d} are same or not may depend on whether $\tilde{\mathcal{M}}_1(K)$ is the same or not as $\tilde{\mathcal{M}}_2(B_d)$. We know the experimental result as

$$\theta_K = (6.5 \pm 0.2) \times 10^{-3}, \quad (54)$$

which is appeared in Ref.[41]. Therefore if this FB-model is admissible, the indirect CP violations of other mesons are also very small and difficult to observe. But as Gérard and Nakada[41] and Soares and Wolfenstein[51] have pointed out, the measurements of asymmetries of $B \rightarrow \psi K$ and 2π decays will distinguish the standard CKM-model from the SW-model. For the same purpose we wish to carry out the precise measurements of ΔM_D ; the dilepton charge asymmetry and also the total charge asymmetry of D^0 - $\overline{D}^0$ system, which surely discriminate which model is true one. If the future experiments confirm that $\Delta M_D \simeq 10^{-14} \sim 10^{-13}$ GeV and $\theta_K \simeq \theta_D \simeq \theta_{B_s}$, it could be said that the physics in TeV energy region (in which **y**-particles play the essential role) totally controls ΔM_P and the indirect CP violations.

Acknowledgements

We would like to thank to the hospitality at the laboratory of the elementary particle physics in Nagoya University.

References

- [1] J.H.Christensen, J.L.Fitch and R.Turlay, Phys. Rev. Lett. **13**(1964)138.
- [2] L.Wolfenstein, Phys. Rev. Lett. **13** (1964)562.
- [3] L.L.Chau. Phys. Rep. **95**(1983)1; A.Pascos and U.Türke, Phys. Rep. **178** (1989)145; P.J.Franzini, Phys. Rep. **173** (1989)1.
- [4] Partice and Fields, Phys. Rev. **D54**(1996)1.
- [5] T.Matsushima, Nuovo Cim. **106A**(1993)139.
- [6] S.Kobayashi, Can. J. Math. **8**(1956)145; Ann. di Math. **43**(1957)119.

- [7] W.Drechsler, J. Math. Phys. **18**(1977)1358; Lecture Notes in Physics,**67**, edited by J.Ehles, et. al. (Springer Verlag)p1.
- [8] R.Gilmore, Lie Groups, Lie Algebras, and Some of Their Applications(Jhon Wiley & Sons 1974) p50.
- [9] H.Geogi and S.L.Glashow, Phys. Rev. Lett. **32**(1974)438.
- [10] J.D.Bjorken, Phys. Rev. **D19**(1979)335.
- [11] P.Q.Hung and J.J.Sakurai, Nucl. Phys. **B143**(1978)81.
- [12] W.Królikowski, Act. Phys. Pol. **B11**(1980)431.
- [13] E.G.H.Harari and N.Seiberg, Phys. Lett. **98B**(1981)269.
- [14] D.W.Greenberg and J.Sucher, Phys. Lett. **99B**(1981)239.
- [15] L. Abbot and E.Farhi, Phys. Lett. **101B** (1981)69.
- [16] H.Fritzsch and G.Mandelbaum, Phys. Lett. **102B**(1981)319; *ibid*,**109B**(1982)224.
- [17] B.Schrempp and F.Schrempp, Nucl. Phys. **B231**(1984)109.
- [18] H.Fritzsch, Lectures given at the International School on Subnuclear Physics, Erice, Sicily August(1984), Max-Plank-Inst. preprint MPI-PAE/PTh85/84 November(1984).
- [19] J.C.Pati and A.Salam, Nucl. Phys. **B214**(1983)109.
- [20] Y.Ohmuki, Unitary Representation of the Poincaré Group and Relativistic Wave Equations(World Scientific, Singapore, 1988)p.1.
- [21] Z.Maki, Proc. of 1981 INS Symposium on Quark and Lepton Physics, edited. by K.Fujikawa et al. p325.
- [22] G.'tHooft, Recent Developments in Gauge Theories,
- [23] F.Abe, et al., Phys. Rev. Lett. **77**(1996)438.
- [24] H.Harari and N.Seiberg Nucl.Phys. **B204**(1982)141.
- [25] L.F.Abbot and E.Farhi, Nucl. Phys. **B189**(1981)547.

- [26] S.Weingerg, Phys. Lett. **102B**(1981)401.
- [27] H.Fritzsch, Lectures in Proc. Arctic School of Physics, Åkä slompolo. Finland (1982).
- [28] D.Schildknecht, Electro weak Effects at High Energies edited by H.B.Newman (Plenum Pub. Comp. 1985)p.1.
- [29] G.Gounaris, R.Kögerler and D.Schildknecht, Phys. Lett. **133B**(1983)118.
- [30] M.K.Gaillard and B.W.Lee, Phys. Rev. **D10** (1974)897.
- [31] L.Wolfenstein, Nucl. Phys. **B160**(1979)501.
- [32] A.Datta and D.Kumbhakar, Z. Phys. **C27** (1985)515.
- [33] A.Datta, Phys. Lett. **B154**(1985)287.
- [34] A.D.Hadeed and B.Holdom, Phys. Lett. **B159**(1985)379.
- [35] I.Picek, Phys. Lett. **B159**(1985)385.
- [36] L.Wolfenstein, Phys. Lett. **B164**(1985)170.
- [37] J.F.Donoghue, E.Golowich, B.R.Holstein and J.Trampetić, Phys. Rev. **D33**(1986)179.
- [38] H.Geogi, Phys. Lett. **B279**(1992)353; T.Ohl, G.Ricciardi and E.H.Simmons, Nucl. Phys. **B403**(1993)605.
- [39] A.Datta and J.C.Pati, Phys. Lett. **B164** (1985)173.
- [40] J.Liu and L.Wolfenstein, Phys. Lett. **B197**(1987)536.
- [41] J.M.Gérard and T.Nakada, Phys. Lett. **B261**(1991)474.
- [42] A.Ali and D.London, Z. Phys. **C65**(1995)431.
- [43] S.L.Wu, CERN-PPE/96-82(1996).
- [44] T.P.Cheng and M.Sher, Phys. Rev. **35** (1987)3484.
- [45] T.Nakada, PSI-PR-91-02 January(1991).

- [46] A.DeRújula, H.Georgi and S.L.Glashow, Phys. Rev. **D12**(1975)147; C.Quigg and J.D.Rosner, Phys. Rep. **56**(1979)167.
- [47] Y.Yamanaka, H.Sugawara and S.Pakvasa, Phys. Rev. **D25**(1982)1895.
- [48] N.Cabibbo, Phys. Rev. Lett. **10**(1963)531; M.Kobayashi and K.Maskawa, Prog. Theor. Phys. **49**(1973)652.
- [49] L.Lavoura, Int. Jour. of Mod. Phys. **A9** (1994)1873.
- [50] G.C.Branco and A.I.Sanda, Phys. Rev. **D26** 1982)3176.
- [51] J.M.Soares and L.Wolfenstein, Phys. Rev. **D47**(1993)1021.
- [52] T.Matsusuima, Quarks, Leptons as Fermon-Boson Composite Objects and Flavor-Mixings by Substructre dynamics TMI-96-2 November(1996).

Fig.(1)

Subquark line-diagrams of the weak interactions




Fig.(2)

Schematic illustrations of P^0 - $\overline{P}^0$ mixings by $\mathbf{y}$ -exchange interactions

